UNIT #5– EXPONENTIAL AND LOGARITHMIC FUNCTIONS MATH III HONORS

Part I Questions

1. The expression $\left(\frac{1}{x^3}\right)^2$ is equivalent to

(1) x^{-1} (3) x^{-5}

- (2) $x^{\frac{2}{3}}$ (4) x^{-6}
- 2. The exponential function $y = 16(2^x)$ could be rewritten as
 - (1) $y = 32^x$ (3) $y = 2^{x+4}$
 - (2) $y = 2^{5x}$ (4) $y = 2^{x^3}$
- 3. The expression $a^{\frac{5}{2}}$ is equivalent to which of the following as long as a > 0?
 - (1) $\sqrt{a^5}$ (3) $\sqrt[5]{a^2}$
 - (2) $\sqrt{5a}$ (4) $\frac{5a}{2}$
- 4. Which of the following would give the same result as $\left(\sqrt[3]{2}\right)^4$?
 - (1) $\sqrt[5]{8}$ (3) $\sqrt{2}$
 - (2) $\sqrt[4]{8}$ (4) $\sqrt[3]{4}$
- 5. For the function $f(x) = 5(2)^{x} + 7$, which of the following represents its y-intercept?
 - (1) 7 (3) 12
 - (2) 5 (4) 17

6. Which of the following could be the equation of the graph shown below?



7. Which of the following values of *x* solves: $(0.5)^{3x+2} = 8^{5x-4}$?

(1) $\frac{2}{3}$	(3) 3
(2) $\frac{5}{9}$	(4) 7

- 8. A population of fruit flies is increasing at a rate of 22.5% per hour. If the population had an original size of 10 flies, then which of the following is its size after one day?
 - (1) 798 (3) 1122
 - (2) 935 (4) 1304

9. The water level in a draining reservoir is changing such that the depth of water decreases by 7.5% per hour. If the water starts at a depth of 45 feet, then which of the following functions properly models the depth, d, as a function of time, t, in hours since it started draining?

- (1) $d = 45(.075)^t$ (3) $d = 45(7.5)^t$
- (2) $d = 45(.925)^t$ (4) $d = 45(92.5)^t$
- 10. The temperature of a cooling liquid in a room held at a constant 75 degrees Fahrenheit can be described by the equation $F(t) = 132(.97)^t + 75$, where *F* is the Fahrenheit temperature and *t* is the amount of time it has been cooling, in minutes. Which of the following was the original temperature of the liquid when it began cooling?
 - (1) 75 (3) 203
 - (2) 132 (4) 207

- 11. If a population grows at a constant rate of 2.8% per year, then by what percent will it grow over the next 10 years?
 - (1) 17% (3) 32%
 - (2) 28% (4) 39%
- 12. The half-life of a radioactive material is the amount of time it takes for 50% of its radioactivity to decrease. If a particular material has a half-life of 35 years, then what percent will remain radioactive after 100 years?
 - (1) 13.8% (3) 34.8%
 - (2) 22.7% (4) 48.7%
- 13. Which of the following is closest to the value of $\log_4(40)$?
 - (1) 1.8 (3) 2.7
 - (2) 2.3 (4) 3.5
- 14. If b > 0 then $\log_b \left(\frac{1}{b^3}\right)$ is equal to (1) $\frac{1}{3}$ (3) 3 (2) $\frac{b}{3}$ (4) -3
- 15. Given the function $f(x) = \log_2(2x-8)$, which of the following values of x is *not* in the domain of the function?
 - (1) x = 5 (3) x = 8
 - (2) x = 2 (4) x = 20

16. Which of the following equations is shown graphed on the grid below?



- (2) $\frac{2\log x}{\sqrt{\log y}}$ (4) $\frac{4\log x \log y}{2}$
- 18. If $\log_b(5) = 1.2$ then $\log_b(125) = ?$
 - (1) 0.4 (3) 3.6
 - (2) 1.728 (4) 30

19. If $5b^{x-3} = 7$ then x =

$(1) \ \frac{\log_b\left(7\right)}{5} + 3$	(3) $3 + \log_b (1.4)$
(2) $\frac{5b}{7} - 3$	(4) $3b^{\frac{7}{5}}$

20. If $f(x) = 50(0.92)^{x} + 75$ then which of the following values of x solves the equation f(x) = 90?

- (1) 12.1 (3) 15.8
- (2) 14.4 (4) 18.3

21. If $ae^{kt} - c = 0$ then which of the following is the value of t based on a, k, and c and the natural base e?

(1)
$$\frac{1}{k} \ln\left(\frac{c}{a}\right)$$
 (3) $\ln\left(\frac{c}{ak}\right)$

(2)
$$\frac{\ln(c)}{ak}$$
 (4) $\frac{ac}{ke}$

- 22. If \$500 is placed in a savings account that earns a 6% nominal interest compounded monthly, then which of the following represents the account's worth after 10 years?
 - (1) \$800.00 (3) \$895.42
 - (2) \$873.29 (4) \$909.70
- 23. How many years, to the nearest tenth, would it take for an investment to double if it is earning a continuous compound interest of 3.5% per year?
 - (1) 17.4 years (3) 22.5 years
 - (2) 19.8 years (4) 25.1 years

24. If a liquid is cooling down according to the formula $y = 84e^{kt} + 55$ and at t = 22 the temperature is y = 71 then which of the following is the value of *k* to the nearest hundredth?

- (1) -0.08 (3) 0.29
- (2) -0.27 (4) 0.58
- 25. The temperature of a cooling liquid is given by the function $T(m) = 38(0.82)^m + 21$, where *T* represents the temperature in degrees Celsius and *m* represents the number of minutes, $m \ge 0$, that the liquid has been cooling. Which of the following represents a temperature that the liquid does not reach as it cools down?
 - (1) 53 (3) 41
 - (2) 16 (4) 28

Free Response Questions

- 26. On the grid shown below, the graph of $f(x) = 2^x$ is shown.
 - (a) On the same graph grid, create an accurate sketch of this function's inverse, $f^{-1}(x)$.

- (b) State the equation of $f^{-1}(x)$.
- (c) State the domain and range of both f(x) and $f^{-1}(x)$.
 - f(x)

Domain:

Range:

27. The expression $(\sqrt[3]{b})^5 \left(\frac{1}{b^2}\right)$ can be written as b^a in simplest form. Determine the value of *a*. Show how you arrived at your answer.

28. If
$$g(x) = \left(\frac{1}{5}\right)^{2x+7} - 3$$
 then algebraically determine the solution to the equation $g(x) = 22$.

Domain:

Range:



29. For the logarithmic function $f(x) = \log_2(x-4)$, explain why x = 0 is not in its domain.

30. For some base, *b*, it is known that $\log_b(5) = 1.28$ and $\log_b(2) = 0.55$. For the same base, determine the value of $\log_b(40)$. Explain how you found your answer.

31. A bank account's worth can be modeled using the formula $w(t) = 380 \left(1 + \frac{.02}{4}\right)^{4t}$, where *w* represents the worth in dollars and *t* represents the number of years since the principal was deposited into the account. Algebraically determine the number of years, to the nearest quarter of a year, it takes for the account to be worth \$500.

Why does it make sense to round your answer to the nearest quarter of a year?

32. If the population of Ashmore, Illinois is decreasing by 5.8% per year, then by what percent will it decrease in the next 5 years? Show how you arrived at your result. Round to the nearest tenth of a percent.

33. A liquid with an initial temperature of 194 °F cools in a room whose temperature is held at 68 °F. The temperature of the liquid, *T*, as it cools can be modeled as a function of time, *x*, using:

$$T(x) = (T_i - T_r)e^{kx} + T_r$$

Where T_i is the initial temperature, T_r is the temperature of the room and k is the decay constant.

(a) If T(15) = 102 then find the value of k accurate to the nearest hundredth.

(b) How many minutes does the model predict it will take for the liquid to reach a temperature of 70 °F? Round to the nearest minute and show or explain how you arrived at your answer.